arXiv:quant-ph/0404174v2 8 Jul 2004

Comment on "Geometric phases for mixed states during cyclic evolutions"

Erik Sjöqvist‡

Department of Quantum Chemistry, Uppsala University, Box 518, Se-751 20 Uppsala, Sweden

Abstract. It is shown that a recently suggested concept of mixed state geometric phase in cyclic evolutions [2004 *J. Phys. A* **37** 3699] is gauge dependent.

PACS numbers: 03.65.Vf, 03.67.Lx

---

‡ Electronic address: erik.sjoqvist@kvac.uu.se

Fu and Chen [1] have proposed a concept of geometric phase for mixed quantal states in cyclic unitary evolution. In the present paper, we demonstrate that this phase is gauge dependent and thus not a property of the path in state space.

Consider the unitary path

$$\mathcal{C} : t \in [0,\tau] \rightarrow \rho(t) = U(t)\rho(0)U^{\dagger}(t) \tag{1}$$

of the density operator $\rho(0)$. The evolution is cyclic iff $[\rho(0), U(\tau)] = 0$. In such a case, $\mathcal{C}$ is closed and the mixed state phase ϕ_g proposed in Ref. [1] takes the form

$$\phi_g = \sum_k w_k \phi_g^k, \tag{2}$$

where w_k are the time-independent eigenvalues of the density operator and

$$\phi_g^k = i \int_0^{\tau} \langle \psi_k(t) | d\psi_k(t) \rangle \tag{3}$$

are the cyclic pure state geometric phases [2] associated with the corresponding eigenvectors $|\psi_k(t)\rangle$ chosen to be periodic, i.e., $|\psi_k(\tau)\rangle = |\psi_k(0)\rangle$, $\forall k$.

Now, the periodicity of the eigenvectors is preserved under the transformation $|\psi_k(t)\rangle \rightarrow |\tilde{\psi}_k(t)\rangle = e^{-i\alpha_k(t)}|\psi_k(t)\rangle$ if $\alpha_k(\tau) - \alpha_k(0) = 2\pi n_k$, n_k integer. It follows from Eq. (3) that

$$\phi_g^k \rightarrow \tilde{\phi}_g^k = i \int_0^{\tau} \langle \tilde{\psi}_k(t) | d\tilde{\psi}_k(t) \rangle = \phi_g^k + 2\pi n_k, \tag{4}$$

i.e., $e^{i\tilde{\phi}_g^k} = e^{i\phi_g^k}$. On the other hand, by inserting Eq. (4) into Eq. (3), we obtain

$$\phi_g \rightarrow \tilde{\phi}_g = \sum_k w_k \tilde{\phi}_g^k = \phi_g + 2\pi \sum_k w_k n_k, \tag{5}$$

where the additional term on the right-hand side is in general not an integer multiple of 2π, i.e., $e^{i\tilde{\phi}_g} \neq e^{i\phi_g}$.

Notice that the path $\mathcal{C}$ is invariant under the gauge transformation $U(t) \rightarrow U(t)V(t)$ if $[\rho(0), V(t)] = 0,\ \forall t \in [0,\tau]$. Explicitly, we may take

$$V(t) = \sum_k e^{-i\alpha_k(t)} |\psi_k(0)\rangle\langle\psi_k(0)| \tag{6}$$

with $\alpha_k(t)$ given above. Due to its dependence upon $V(t)$ via the integers n_k, it follows that ϕ_g is not a property of the closed path $\mathcal{C}$. In other words, the phase concept proposed in Ref. [1] is not gauge invariant.

A gauge invariant mixed state geometric phase has been proposed in Ref. [3]. In the cyclic case, this phase γ is determined by

$$\mathcal{V} e^{i\gamma} = \sum_k w_k e^{i\phi_k^g} \tag{7}$$

and is independent of n_k as each phase factor in the sum of the right-hand side transforms invariantly. This phase may be tested interferometrically as an incoherent average of pure state interference profiles upon elimination of all the pure state dynamical phases in one of the arms. For cyclic evolution, such an analysis yields the intensity

$$\mathcal{I} \propto \sum_k w_k \Big| e^{i\chi} |\psi_k(0)\rangle + e^{i\phi_g^k} |\psi_k(0)\rangle \Big|^2 \propto 1 + \mathcal{V} \cos\left[\chi - \gamma\right], \tag{8}$$

where χ is a variable U(1) shift applied to the other arm. γ has recently been measured using nuclear magnetic resonance technique [4].

In conclusion, we have demonstrated that a quantal phase concept recently proposed in Ref. [1] is gauge dependent and therefore not a property of the path in state space. Thus, this phase is neither experimentally testable, nor does it qualify as a geometric phase for mixed states during cyclic evolutions.

References

[1] Fu L-B and Chen J-L 2004 *J. Phys. A* **37** 3699

[2] Aharonov Y and Anandan J 1987 *Phys. Rev. Lett.* **58** 1593

[3] Sjöqvist E, Pati A K, Ekert A, Anandan J S, Ericsson M, Oi D K L, and Vedral V 2000 *Phys. Rev. Lett.* **85** 2845

[4] Du J F, Zou P, Shi M, Kwek L C, Pan J-W, Oh C H, Ekert A, Oi D K L, and Ericsson M 2003 *Phys. Rev. Lett.* **91** 100403